

## P V and the Mass Loss Discrepancy in O Stars

D. Massa

*SGT, Inc.*

*Code 681, NASA's GSFC, Greenbelt, MD 20901, USA*

A.W. Fullerton

*University of Victoria & Johns Hopkins University*

*3400 N. Charles St., Baltimore, MD 21218, USA*

R.K. Prinja

*Department of Physics & Astronomy, UCL*

*Gower Street, London WC1E 6BT, UK*

**Abstract.** Building upon a previous analysis of P v wind lines in LMC O stars, we analyze the P v wind lines in a sample of Galactic O stars which have empirical mass loss rates determined from either their radio fluxes or H $\alpha$  profiles. Since the wind analysis provides a measure of  $\dot{M}q$  where  $q$  is the ionization fraction of the ion, we determine  $q(\text{P v})$  observationally. In spite of model predictions that  $q \sim 1$  for mid-O stars, we find  $q(\text{P v}) \leq 0.15$  throughout the O stars. We discuss the origin of this discrepancy.

### 1. Measures of $\dot{M}$

Three approaches are normally used to determine stellar mass loss rates,  $\dot{M}$ s. All assume that the wind is homogeneous and spherically symmetric (SS) with a single, monotonic velocity law, and all *should* agree. The three approaches are:  
*1. Continuum excess from free-free emission.* This samples the outer wind (the exact radius depends on wavelength), where it becomes optically thick to free-free emission. It is only detectable for massive winds in nearby stars. The radio wavelengths are considered “cleanest”, because, in contrast to the IR/FIR, massive winds become optically thick at large radii ( $\geq 10R_*$ ), where  $v = v_\infty$ , a constant. This makes  $\rho_{\text{wind}} \sim \dot{M}/(r^2 v_\infty)$ , independent of  $v(r)$ . Furthermore, no photospheric correction is needed. However, the radio flux can be non-thermal, so observations at multiple wavelengths are required to determine the spectral index of the emission.

*2. H $\alpha$  emission.* This samples the inner wind and is easily observed. For massive winds, H $\alpha$  emission is related to  $\dot{M}$ . However, the exact form of the observed H $\alpha$  profile depends upon the  $N = 3$  departure coefficient for H in the wind and this, in turn, depends upon: the photospheric radiation field; the diffuse radiation field of the wind; and the wind velocity law in the acceleration region. The shape of the “photospheric” H $\alpha$  profile is also required, and the observed  $W_\lambda(\text{H}\alpha)$  can be strongly variable. Nevertheless, relatively sophisticated models

for  $H\alpha$  formation exist (e.g., Repolust et al. 2004), and can provide reasonable agreement between available radio and  $H\alpha$   $\dot{M}$ s when  $H\alpha$  emission is strong.

3. *UV resonance lines.* These sample the entire wind. Their shapes are determined by the radial optical depth of the wind,  $\tau_{rad} \sim \dot{M} q_i A_E$ , where  $A_E$  and  $q_i$  are the abundance of element  $E$  and its ionization fraction for stage  $i$ . However observations of a dominant ion ( $q_i \sim 1$ ), of known abundance are required to estimate  $\dot{M}$  directly, and the wind lines of abundant, dominant ions are saturated in winds massive enough to be detected in the radio or to have reliable  $H\alpha$   $\dot{M}$ s.

## 2. FUSE and P V

*FUSE* gives access to P v  $\lambda\lambda 1118, 1128$ . P v is a surrogate for C iv (Massa et al. 2003) and  $q_i \sim 1$  is expected for both ions in mid-O star winds. Unlike  $A_C$ ,  $A_P \simeq Const$  over the life of an O star. Furthermore, for scaled solar abundances,  $\tau_{rad}(C\text{ iv})/\tau_{rad}(P\text{ v}) = 661$ , so to detect P v,  $\tau_{rad}(C\text{ iv}) \geq 50$  and saturated – as is the case for stars with detectable radio fluxes and strong  $H\alpha$  emission.

**P V in LMC O Stars:** The first large scale *FUSE* study of P v was by Massa et al. (2003). They performed SEI (Lamers et al. 1987) fits to P v wind lines, determining  $\tau_{rad}$  for 25 LMC O stars. They then used  $\dot{M}$ s predicted by the Vink et al. (2001) theory and  $A_P = 0.50 \times \text{solar}$  to find that  $q(P\text{ V})$  peaked between 45–50 kK, as expected, but with a peak value  $\leq 0.15$  (see Fig. 1). This result implies a factor of 7 or more discrepancy between the expected and observed  $\dot{M}$ s. There are three possible explanations for this result:

1. The LMC P abundance scales differently from other elements,
2. The theoretical  $\dot{M}$ s are incorrect for the LMC, or
3. The winds are not homogeneous and SS, but strongly clumped or structured.

**P V in Galactic O Stars (preliminary results):** The Galactic  $A_P$  is well determined (e.g., Catanzaro et al. 2003), so abundance is not an issue for the Galaxy. We are currently analyzing P v in *Copernicus*, *Orfeus* and *FUSE* data for stars with radio and/or  $H\alpha$   $\dot{M}$  estimates. This eliminates the need for model  $\dot{M}$ s to derive  $q$  for P v, and determines how well different mass loss rate indicators agree. So far, we have analyzed 30 stars (11 more will be observed by *FUSE*), and the results are shown in Figure 2. These results are effectively identical to those from the LMC (Fig. 1), implying the same conclusions.

## 3. Conclusions

- An erroneous P abundance is not the cause of the small LMC  $q(P\text{ V})$ s.
- Strong clumping/porosity must be the root of the problem.
- Large scale clumping can also strongly affect the radio and  $H\alpha$   $\dot{M}$ s, so these should be re-evaluated.
- The good news: the different measures of  $\dot{M}$  are sensitive to different aspects of the wind flow, so bringing them all into agreement (together with the X-ray and O vi wind line observations) will provide powerful constraints on models of how the winds are structured.




Figure 1. Mean P v ionization fractions as a function of temperature for LMC O stars analyzed by Massa et al. (2003). Open, half filled and filled symbols denote stellar luminosities in the ranges  $\log L/L_{\odot} > 6.0$ ,  $6.0 \geq \log L/L_{\odot} > 5.6$  and  $5.6 \geq \log L/L_{\odot}$ , respectively.




Figure 2. Mean P v ionization fractions versus temperature for Galactic O stars with radio or  $H\alpha$   $\dot{M}$ s. Filled symbols: radio  $\dot{M}$ s, open circles:  $H\alpha$   $\dot{M}$ s from Repolust et al. (2004), open triangles:  $H\alpha$   $\dot{M}$ s from Lamers & Leitherer (1993).

## References

- Catanzaro, G., André, M.K., Leone, F. & Sonnentrucker, P. 2003 A&A, 404, 667  
 Lamers, H.J.G.L.M., Cerruti-Sola, M., & Perinotto, M. 1987, ApJ, 314, 726  
 Lamers, H.J.G.L.M. & Leitherer, C. 1993 ApJ, 412, 771  
 Massa, D., Fullerton, A.W., Sonneborn, G. & Hutchings, J.B. 2003 ApJ, 586, 996  
 Repolust, T., Puls, J., & Herrero, A. 2004 A&A, 415, 349  
 Vink, J.S., de Koter, A., & Lamers, H.J.G.L.M. 2001, A&A, 369, 574